

## Effect of Endophytic Fungus, *Piriformospora Indica*, on Barley Resistance to Lead

F. Karimi<sup>\*1</sup>, M. Sepehri<sup>2</sup>, M. Afuni<sup>3</sup> and M. A. Hajabbasi<sup>4</sup>

(Received: Nov. 27-2011 ; Accepted : Oct. 21-2014)

### Abstract

By modifying plants at genetical, physiological and ecological levels, entophytic fungi as the most important soil microorganisms have a pronounced growth-promoting activity and also increase plant resistance to biotic and abiotic stresses. This research was undertaken to evaluate the potential of *P. indica* to increase barley (*Hordeumvulgare* L.) resistance to lead (pb). Therefore, a greenhouse experiment with two fungus treatments (non-inoculated and *P. indica* inoculated) and five levels of pb (0, 25, 50, 100 and 500 mg/kg) with three replications was conducted based on a factorial design. Measurement of shoot and root dry weight showed that the growth of *P. indica*-colonized plants at all levels of pb treatments was higher ( $P < 0.05$ ) than that of the corresponding controls. Also, chlorophyll concentration of inoculated plants with *P. indica* was superior to non-inoculated plants. In addition, the results showed that in contrast to the plant shoot, lead concentration in the root of *P. indica*-colonized plants was higher than the non-inoculated controls.

**Keyword:** *Piriformospora indica*, Lead, Barley.

---

1. Dept. of Soil Sci., College of Agric., Isf. Univ. Technol., Isfahan, Iran.

\*: Corresponding Author, Email: fatemeh.karimi@rocketmail.com